www.innocyto.com

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

iF488 Anti-human/mouse/rat CD278 (ICOS) Antibody

Catalog Number:	111205, 111206
Size:	25 tests, 100 tests
Target Name:	CD278, ICOS, Inducible Costimulatory molecule
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	C398.4A
Application:	Flow Cytometry
Reactivity:	Human, mouse, rat
Format:	iF488
Isotype:	Armenian Hamster IgG
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 ÂµL of this reagent per 0.5-1.0 million cells in a 100 ÂµL volume. Optimal reagent performance should be determined by titration for each specific application. iF488 has an excitation max at 491 nm and an emission max at 516 nm.
Excitation Laser:	Blue Laser (488 nm)
Isotype Control:	300507

BACKGROUND INFORMATION

ICOS is a 47–57 kDa costimulatory molecule homologous to CD28/CTLA-4 and is expressed on activated T cells and a subset of thymocytes. Engagement of ICOS enhances T cell proliferation, promotes cytokine production such as IL-4 and IL-10, and plays an important role in humoral immunity and Th2 cell differentiation.

PRODUCT DATA



Human peripheral blood lymphocytes stimulated with plate-coated anti-Human CD3 and CD28 for three days were stained with iF488 Anti-Human/Mouse/Rat ICOS clone C398.4A (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.