

Anti-Mouse CD16.2 (FcγRIV) Antibody

Catalog Number:	205201, 205202
Size:	25 µg, 100 µg
Target Name:	CD16.2, Fc gamma RIV, FcγRIV, Fcγr4, FcγgammaRIV
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	9E9
Application:	Flow Cytometry
Reactivity:	Mouse
Format:	Purified
Isotype:	Armenian Hamster IgG
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Protein Concentration:	0.5 mg/mL
Storage&Handling:	The antibody solution should be stored between 2°C and 8°C
Isotype Control:	300501

BACKGROUND INFORMATION

Mouse CD16.2 (FcγRIV) is a low-affinity activating Fc gamma receptor expressed on myeloid immune cells, including monocytes, macrophages, neutrophils, and certain dendritic cell subsets. It belongs to the immunoglobulin superfamily and mediates effector functions triggered by IgG immune complexes.

It is a type I transmembrane glycoprotein with an extracellular Ig-like domain responsible for binding IgG, a single-pass transmembrane region, and a short cytoplasmic tail. Unlike inhibitory Fcγ receptors, CD16.2 signals through association with the common Fc receptor gamma chain, which contains immunoreceptor tyrosine-based activation motifs (ITAMs) that initiate downstream signaling.

CD16.2 primarily binds IgG2a and IgG2b immune complexes in mice, promoting antibody-dependent cellular cytotoxicity, phagocytosis, and cytokine release. It plays a central role in host defense against pathogens but can also contribute to inflammatory and autoimmune pathology when overactivated.

In therapeutic contexts, CD16.2 is important for monoclonal antibody efficacy in preclinical mouse models, as its engagement enhances ADCC and tumor clearance. Conversely, blocking or modulating FcγRIV signaling is being explored to reduce autoimmune disease severity and excessive inflammation.

Overall CD16.2 serves as a key activating Fc receptor that bridges humoral immunity and cellular effector responses. Its balanced signaling determines whether IgG complexes drive protective immunity or pathological inflammation. Because of its strong role in antibody effector functions, it is widely used in mouse models to evaluate therapeutic antibodies and engineer improved Fc designs

for enhanced clinical performance in vivo studies.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.